

Rapid Note

Inverse cubic law for the distribution of stock price variations

P. Gopikrishnan^a, M. Meyer, L.A.N. Amaral, and H.E. Stanley

Center for Polymer Studies and Department of Physics, Boston University Boston, MA 02215, USA

Received: 23 April 1998 / Revised and Accepted: 24 April 1998

Abstract. The probability distribution of stock price changes is studied by analyzing a database (the Trades and Quotes Database) documenting every trade for all stocks in three major US stock markets, for the two year period January 1994 – December 1995. A sample of 40 million data points is extracted, which is substantially larger than studied hitherto. We find an asymptotic power-law behavior for the cumulative distribution with an exponent $\alpha \approx 3$, well outside the Lévy regime ($0 < \alpha < 2$).

PACS. 89.90.+n Other areas of general interest to physicists

The asymptotic behavior of the increment distribution of economic indices has long been a topic of avid interest [1–6]. Conclusive empirical results are, however, difficult to obtain, since the asymptotic behavior can be obtained only by a proper sampling of the tails, which requires a huge quantity of data. Here, we analyze a database documenting each and every trade in the three major US stock markets, the New York Stock Exchange (NYSE), the American Stock Exchange (AMEX), and the National Association of Securities Dealers Automated Quotation (NASDAQ) for the entire 2-year period January 1994 to December 1995 [7]. We thereby extract a sample of approximately 4×10^7 data points, which is much larger than studied hitherto.

We form 1000 time series $S_i(t)$, where S_i is the market price of company i (*i.e.* the share price multiplied with the number of outstanding shares), $i = 1 \dots 1000$ is the rank of the company according to its market price on January 1, 1994. The basic quantity of our study is the relative price change,

$$G_i(t) \equiv \ln S_i(t + \Delta t) - \ln S_i(t) \simeq \frac{S_i(t + \Delta t) - S_i(t)}{S_i(t)}, \quad (1)$$

where the time lag is $\Delta t = 5$ min. We normalize the increments,

$$g_i(t) \equiv [G_i(t) - \langle G_i(t) \rangle] / v_i, \quad (2)$$

where the volatility $v_i \equiv \sqrt{\langle (G_i(t) - \langle G_i(t) \rangle)^2 \rangle}$ of company i is measured by the standard deviation, and $\langle \dots \rangle$ is a time average [8].

We obtain about 20 000 normalized increments $g_i(t)$ per company per year, which gives about 4×10^7 increments for the 1000 largest companies in the time period

studied. Figure 1a shows the cumulative probability distribution, *i.e.* the probability for an increment larger or equal to a threshold g , $P(g) \equiv P\{g_i(t) \geq g\}$, as a function of g . The data are well fit by a power law

$$P(g) \sim g^{-\alpha} \quad (3)$$

with exponents $\alpha = 3.1 \pm 0.03$ (positive tail) and $\alpha = 2.84 \pm 0.12$ (negative tail) from two to hundred standard deviations.

In order to test this result, we calculate the inverse of the local logarithmic slope of $P(g)$, $\gamma^{-1}(g) \equiv -d \log P(g) / d \log g$ [9]. We estimate the asymptotic slope α by extrapolating γ as a function of $1/g$ to 0. Figure 1b shows the results for the negative and positive tail respectively, each using all increments larger than 5 standard deviations. Extrapolation of the linear regression lines yield $\alpha = 2.84 \pm 0.12$ for the positive and $\alpha = 2.73 \pm 0.13$ for the negative tail. We test this method by analyzing two surrogate data sets with known asymptotic behavior, (a) an independent random variable with $P(x) = (1+x)^{-3}$ and (b) an independent random variable with $P(x) = \exp(-x)$. The method yields the correct results 3 and ∞ respectively.

To test the robustness of the inverse cubic law $\alpha \approx 3$, we perform several additional calculations: (*i*) we change the time increment in steps of 5 min up to 120 min, (*ii*) we analyze the S&P500 index over the 13y-period January '84 – December '96 using the same methods as above (Fig. 1c and d), (*iii*) we replace definition of the volatility by other measures, such as $v_i \equiv \langle |G_i(t) - \langle G_i(t) \rangle| \rangle$. The results are all consistent with $\alpha = 3$. These extensions will be discussed in detail elsewhere [10].

To put these results in the context of previous work, we recall that proposals for $P(g)$ have included (*i*) a Gaussian distribution [1], (*ii*) a Lévy distribution [2, 11, 12], and

^a e-mail: gopi@bu.edu




Fig. 1. (a) Log-log plot of the cumulative probability distribution $P(g)$ of the normalized price increments $g_i(t)$. The lines in (a) are power law fits to the data over the range $2 < g < 100$. (b) The inverse local slope of $P(g)$, $\gamma(g) \equiv -(d \log P(g)/d \log g)^{-1}$ as a function of $1/g$ for the negative (\circ) and the positive ($+$) tail respectively. We obtain an estimator for $\gamma(g)$, by sorting the normalized increments by their size, $g^{(1)} > g^{(2)} > \dots > g^{(N)}$. The cumulative density can then be written as $P(g^{(k)}) = k/N$, and we obtain for the local slope $\gamma(g^{(k)}) = -\log(g^{(k+1)}/g^{(k)})/\log(P(g^{(k+1)})/P(g^{(k)})) \simeq k(\log(g^{(k+1)}) - \log(g^{(k)}))$. Each data point shown in b is an average over 1000 increments $g^{(k)}$, and the lines are linear regression fits to the data. Note that the average $m^{-1} \sum_{k=1}^m \gamma(g^{(k)})$ over all events used would be identical to the estimator for the asymptotic slope proposed by Hill [9]. (c) Same as (a) for the 1 min S&P500 increments. The regression lines yield $\alpha = 2.93$ and $\alpha = 3.02$ for the positive and negative tails respectively. (d) Same as (b) for the 1 min S&P500 increments, except that the number of increments per data point is 100.

(iii) a truncated Lévy distribution, where the tails become “approximately exponential” [3]. The inverse cubic law differs from all three proposals: Unlike (i) and (iii), it has diverging higher moments (larger than 3), and unlike (i) and (ii) it is not a stable distribution.

We thank X. Gabaix, S. Havlin, Y. Liu, R. Mantegna, C.K. Peng and D. Stauffer for helpful discussions, and JNICT DFG and NSF for financial support.

References

1. L. Bachelier, Ann. Sci. École Norm. Sup. **3**, 21 (1900).
2. B.B. Mandelbrot, J. Business **36**, 294 (1963).
3. R.N. Mantegna, H.E. Stanley, Nature **376**, 46 (1995).
4. S. Ghashgaie, W. Breymann, J. Peinke, P. Talkner, Y. Dodge, Nature **381**, 767 (1996).
5. A. Pagan, J. Empirical Finance **3**, 15 (1996), and references therein.
6. J.P. Bouchaud, M. Potters, *Théorie des Risques Financiers* (Alea-Saclay, Eyrolles 1998); R. Cont, Eur. Phys. J. B (1998), in press (cond-mat/9705075).
7. *The Trades and Quotes Database*, 24 CD-ROM for '94-'95, published by the New York Stock Exchange.
8. In order to obtain uncorrelated estimators for the volatility and the price increment at time t , the time average extends only over time steps $t' \neq t$.
9. B.M. Hill, Ann. Stat. **3**, 1163 (1975).
10. P. Gopikrishnan, M. Meyer, L.A.N. Amaral, H.E. Stanley (to be published).
11. V. Pareto, *Cours d'Économie Politique* (Lausanne and Paris, 1897).
12. P. Lévy, *Théorie de l'addition des variables aléatoires* (Gauthier-Villars, Paris, 1937).