

Universal and Nonuniversal Properties of Cross Correlations in Financial Time Series

Vasiliki Plerou,^{1,2} Parameswaran Gopikrishnan,¹ Bernd Rosenow,³ Luís A. Nunes Amaral,¹ and H. Eugene Stanley¹

¹Center for Polymer Studies and Department of Physics, Boston University, Boston, Massachusetts 02215

²Department of Physics, Boston College, Chestnut Hill, Massachusetts 02167

³Institut für Theoretische Physik, Universität zu Köln, D-50937 Köln, Germany

(Received 22 February 1999)

We use methods of random matrix theory to analyze the cross-correlation matrix \mathbf{C} of stock price changes of the largest 1000 U.S. companies for the 2-year period 1994–1995. We find that the statistics of most of the eigenvalues in the spectrum of \mathbf{C} agree with the predictions of random matrix theory, but there are deviations for a few of the largest eigenvalues. We find that \mathbf{C} has the universal properties of the Gaussian orthogonal ensemble of random matrices. Furthermore, we analyze the eigenvectors of \mathbf{C} through their inverse participation ratio and find eigenvectors with large ratios at both edges of the eigenvalue spectrum—a situation reminiscent of localization theory results.

PACS numbers: 87.23.Ge, 02.50.Ey, 05.40.–a, 05.45.Tp

There has been much recent work applying physics concepts and methods to the study of financial time series [1–15]. In particular, the study of correlations between price changes of different stocks is both of scientific interest and of practical relevance in quantifying the risk of a given stock portfolio [1,2]. Consider, e.g., the equal-time correlation of stock-price changes for a given pair of companies. Since the market conditions may not be stationary, and the historical records are finite, it is not clear if a measured correlation of price changes of two stocks is due only to “noise,” or genuinely arises from the interactions between the two companies. Moreover, unlike most physical systems, there is no “algorithm” to calculate the “interaction strength” between two companies (as there is for, say, two spins in a magnet). The problem is that although every pair of companies should interact either directly or indirectly, the precise nature of interaction is unknown.

In some ways, the problem of interpreting the correlations between individual stock-price changes is reminiscent of the difficulties experienced by physicists in the 1950s, in interpreting the spectra of complex nuclei. Large amounts of spectroscopic data on the energy levels were becoming available but were too complex to be explained by model calculations because the exact nature of the interactions was unknown. Random matrix theory (RMT) was developed in this context to deal with the statistics of energy levels of complex quantum systems [16]. With the minimal assumption of a random Hamiltonian, given by a real symmetric matrix with independent random elements, a series of remarkable predictions was made and successfully tested on the spectra of complex nuclei [16]. RMT predictions represent an average over all possible interactions [16]. Deviations from the *universal* predictions of RMT identify system-specific, nonrandom properties of the system under consideration, providing clues about the nature of the underlying interactions [17,18].

Here, we apply RMT methods to study the cross correlations [11,12] of stock-price changes. First, we demon-

strate the validity of the universal predictions of RMT for the eigenvalue statistics of the cross-correlation matrix. Second, we calculate the deviations of the empirical data from the RMT predictions, obtaining information that enables us to identify cross correlations between stocks not explainable purely by randomness.

We analyze a database [19] containing the price $S_i(t)$ of stock i at time t , where $i = 1, \dots, N$ denotes the N largest U.S. publicly traded companies and the time t runs over the 2-year period 1994–1995. From the time series $S_i(t)$, we calculate the price change $G_i(t, \Delta t)$, defined as

$$G_i(t, \Delta t) \equiv \ln S_i(t + \Delta t) - \ln S_i(t). \quad (1)$$

We report here results for $N = 1000$ and $\Delta t = 30$ min (there are 6448 data points for each of the 1000 companies). The simplest measure of correlations between different stocks is the equal-time cross-correlation matrix \mathbf{C} , which has elements

$$C_{ij} \equiv \frac{\langle G_i G_j \rangle - \langle G_i \rangle \langle G_j \rangle}{\sigma_i \sigma_j}. \quad (2)$$

Here $\sigma_i \equiv \sqrt{\langle G_i^2 \rangle - \langle G_i \rangle^2}$ is the standard deviation of the price changes of company i , and $\langle \dots \rangle$ denotes a time average over the period studied [19].

We analyze the statistical properties of \mathbf{C} by applying RMT techniques. First, we diagonalize \mathbf{C} and obtain its eigenvalues λ_k —with $k = 1, \dots, 1000$ —which we rank from the smallest to the largest. Next, we calculate the eigenvalue distribution [11] and compare it with recent analytical results for a cross-correlation matrix generated from finite uncorrelated time series [20]. Figure 1 shows the eigenvalue distribution of \mathbf{C} , which deviates from the predictions of Ref. [20], for large eigenvalues $\lambda_k \geq 1.94$ (see Fig. 1 caption). This result is in agreement with the results of Ref. [11] for the eigenvalue distribution of \mathbf{C} for $\Delta t = 1$ day.

To test for universal properties, we first calculate the distribution of the nearest-neighbor spacings $s \equiv \lambda_{k+1} - \lambda_k$, which we compute after transforming the eigenvalues in




FIG. 1. The probability density function of the eigenvalues of the normalized cross-correlation matrix C for the 1000 largest U.S. companies in the TAQ database for the 2-year period 1994–1995 [19]. Recent analytical results [20] for cross-correlation matrices generated from uncorrelated time series predict a finite range of eigenvalues depending on the ratio R of the length of the time series to N [11]. In our case $R = 6.448$, corresponding to eigenvalues distributed in the interval $0.37 \leq \lambda_k \leq 1.94$ [20]. However, the largest eigenvalue, $\lambda_{1000} \approx 50$, for the 2-year period (arrow in inset) is ≈ 25 times larger than the maximum eigenvalue predicted for uncorrelated time series. The inset also shows the largest eigenvalue for the four half-year intervals—denoted A, B, C, D . The distribution of eigenvector components for the large eigenvalues, the ones well outside the bulk, show significant deviations from the Gaussian prediction of RMT, which suggests “collective” behavior or correlations [18] between different companies. The largest eigenvalue would then correspond to the correlations within the entire market [11].

such a way that their distribution becomes uniform—a procedure known as unfolding [17,18,21]. Figure 2(a) shows the distribution for the empirical data and compares it with the RMT predictions for real symmetric random matrices. This class of matrices shares universal properties with the ensemble of matrices whose elements are distributed according to a Gaussian probability measure—the Gaussian orthogonal ensemble (GOE). We find good agreement between the empirical data and the GOE prediction,

$$P_{GOE}(s) = \frac{\pi s}{2} \exp\left(-\frac{\pi}{4}s^2\right). \quad (3)$$

A second independent test of the GOE is the distribution of *next-nearest-neighbor* spacings between the rank-ordered eigenvalues [17]. This distribution is expected to be identical to the distribution of nearest-neighbor spacings of the Gaussian symplectic ensemble (GSE), as verified by the empirical data [Fig. 2(b)].

The distribution of eigenvalue spacings reflects correlations only of consecutive eigenvalues but does not contain information about correlations of longer range. To probe any “long-range” correlations, we first calculate the





FIG. 2. Comparison of the RMT predictions for the spacing distributions with results for the empirical cross-correlation matrix. (a) Nearest-neighbor (nn) spacing distribution of the eigenvalues of C after unfolding using the Gaussian broadening procedure [21]. The eigenvalue distribution can be considered to be a sum of delta functions located at each eigenvalue. Each delta function is then “broadened” by choosing a Gaussian distribution with standard deviation $(\lambda_{k+a} - \lambda_{k-a})/2$, where $2a$ is the size of the window used for broadening [21], and the optimum value of a obtained from Fig. 2(d). The solid line is the GOE prediction, Eq. (3), and the dashed line is a fit to the one-parameter Brody distribution $p(s) \equiv B(1 + \beta)s^\beta \exp(-Bs^{\beta+1})$, with $B \equiv [\Gamma(\frac{\beta+2}{\beta+1})]^{1+\beta}$. The fit yields $\beta = 0.99 \pm 0.02$, in good agreement with the GOE prediction $\beta = 1$. At the 80% confidence level, a Kolmogorov-Smirnov test cannot reject the hypothesis that the GOE is the correct description. (b) Next-nearest-neighbor (nnn) spacing distribution of C . RMT predicts that, for the GOE, the distribution of next-nearest-neighbor spacings should follow the same distribution as the nearest-neighbor spacings for the GSE. This prediction is confirmed for the empirical data, both visually and by a Kolmogorov-Smirnov test that at the 40% confidence level cannot reject the hypothesis that the GSE is the correct distribution. (c) Number variance and (d) spectral rigidity of C for different values of the unfolding parameter a , compared to the exact expression for the GOE (solid line) and the uncorrelated case (dashed line). As a increases, both the number variance and the spectral rigidity approach the theoretical curve for the GOE, while the spacing distribution remains essentially unchanged. We choose $a = 15$ as the optimal value.

number variance Σ^2 —the variance of the number of unfolded eigenvalues in intervals of length L around each of the eigenvalues [17,18,21,22]. If the eigenvalues are uncorrelated, $\Sigma^2 \sim L$, while for the case of a “rigid” eigenvalue spectrum, $\Sigma^2 = 0$. For the GOE case, we find the “intermediate” behavior $\Sigma^2 \sim \ln L$, as predicted by RMT [Fig. 2(c)].

A second way to measure long-range correlations in the eigenvalues is through the spectral rigidity Δ —the least square deviation of the unfolded cumulative eigenvalue density from a fit to a straight line in an interval of length L [17,18,21,23]. For uncorrelated eigenvalues, $\Delta \sim L$, whereas for the rigid case $\Delta = \text{const.}$ For the GOE case, we find $\Delta \sim \ln L$ as predicted by RMT [Fig. 2(d)].

Having demonstrated that the eigenvalue statistics of \mathbf{C} satisfy the RMT predictions, we now proceed to analyze the eigenvectors of \mathbf{C} . RMT predicts that the components of the normalized eigenvectors of a GOE matrix are distributed according to a Gaussian probability distribution with mean zero and variance one. In agreement with recent results [11], we find that eigenvectors corresponding to *most* eigenvalues in the “bulk” ($\lambda_k < 2$) follow this prediction. On the other hand, eigenvectors with eigenvalues outside the bulk ($\lambda_k > 2$) show marked deviations from the Gaussian distribution. In particular, the eigenvector corresponding to the largest eigenvalue λ_{1000} deviates significantly from the Gaussian distribution predicted by RMT.

The component ℓ of a given eigenvector relates to the contribution of company ℓ to that eigenvector. Hence, the distribution of the components contains information about the number of companies contributing to a specific eigenvector. In order to distinguish between one eigenvector with approximately equal components and another with a small number of large components, we define the inverse participation ratio [17,24]

$$I_k \equiv \sum_{\ell=1}^{1000} [u_{k\ell}]^4, \quad (4)$$

where $u_{k\ell}$, $\ell = 1, \dots, 1000$ are the components of eigenvector k . The physical meaning of I_k can be illustrated by two limiting cases: (i) an eigenvector with identical components $u_{k\ell} \equiv 1/\sqrt{N}$ has $I_k = 1/N$, whereas (ii) an eigenvector with one component $u_{k1} = 1$ and all the others zero has $I_k = 1$. Therefore, I_k is related to the reciprocal of the number of eigenvector components significantly different from zero.

Figure 3 shows I_k for eigenvectors of a matrix generated from uncorrelated time series with a power-law distribution of price changes [7,8]. The average value of I_k is $\langle I \rangle \approx 3 \times 10^{-3} \approx 1/N$, indicating that the eigenvectors are *extended* [24,25]—i.e., almost all companies contribute to them. Fluctuations around this average value are confined to a narrow range. On the other hand, the empirical data show deviations of I_k from $\langle I \rangle$ for a few of the largest eigenvalues. These I_k take values of the order of 10^{-2} , suggesting that there are groups of approximately




FIG. 3. Inverse participation ratio I_k for each of the 1000 eigenvectors. As a control, we show in the inset the I_k values for the eigenvectors of a cross-correlation matrix computed from uncorrelated independent power-law distributed time series [7,8] of the same length as the data. Empirical data show marked peaks at both edges of the spectrum, whereas the control shows only small fluctuations around the average value $\langle I \rangle = 3 \times 10^{-3}$. The large I_k values for the largest eigenvalues are to be expected from Fig. 1, but the large values of I_k for the small eigenvalues are surprising. We test that the deviations for small eigenvalues are real by comparing our results against controls with different sizes and by analyzing separately each of the four half-year intervals in the database.

50 companies contributing to these eigenvectors. The corresponding eigenvalues are well outside the bulk, suggesting that these companies are correlated [18].

Surprisingly, we also find that there are I_k values as large as 0.27 for eigenvectors corresponding to the smallest eigenvalues $\lambda_i \approx 0.25$ [26]. The deviations from $\langle I \rangle$ for the smallest eigenvalues are about 10^2 to 10^3 times larger than the standard deviation of the fluctuations for the control, which suggests that the eigenvectors are *localized* [24,25]—i.e., only a few companies contribute to them. The small size of the corresponding eigenvalues suggests that these companies are uncorrelated with one another.

The presence of vectors with large I_k also arises in localization theory, where one frequently finds “random band matrices” [24] containing extended states with small I_k in the middle of the band, whereas edge states are localized and have large I_k . Our finding of localized states for small and large eigenvalues of the cross-correlation matrix \mathbf{C} is reminiscent of Anderson localization [27] and suggests that \mathbf{C} may be a random band matrix [28].

In summary, we find that the most eigenvalues in the spectrum of the cross-correlation matrix of stock price changes agree surprisingly well with the *universal* predictions of random matrix theory. In particular, we find that \mathbf{C} satisfies the universal properties of the Gaussian orthogonal ensemble of real symmetric random

matrices. We find through our analysis of the inverse participation ratio of its eigenvectors that \mathbf{C} may be a random band matrix, which may support the idea that a metric can be defined on the space of companies and that a distance can be defined between pairs of companies [29]. Hypothetically, the presence of localized states may allow us to draw conclusions about the “spatial dimension” of the set of stocks studied here, and about the “range” of the correlations between the companies.

We thank M. Barthélémy, N. V. Dohkolyan, X. Gabaix, U. Gerland, S. Havlin, R.N. Mantegna, Y. Lee, C.-K. Peng, D. Stauffer, and H.A. Weidenmüller for helpful discussions, and NSF and FCT/Portugal for financial support.

- [1] *Econophysics: An Emerging Science*, edited by I. Kondor and J. Kertész (Kluwer, Dordrecht, 1999); J.-P. Bouchaud and M. Potters, *Theorie des Risques Financières* (Alea-Saclay, Eyrolles, 1997); R.N. Mantegna and H.E. Stanley, *Introduction to Econophysics: Correlations and Complexity in Finance* (Cambridge University Press, Cambridge, England, 1999).
- [2] E. J. Elton and M. J. Gruber, *Modern Portfolio Theory and Investment Analysis* (J. Wiley, New York, 1995).
- [3] T. Lux and M. Marchesi, *Nature* (London) **297**, 498 (1999).
- [4] J.-P. Bouchaud and R. Cont, *Eur. Phys. J. B* **6**, 543 (1998); M. Potters, R. Cont, and J.-P. Bouchaud, *Europhys. Lett.* **41**, 239 (1998); J.-P. Bouchaud and D. Sornette, *J. Phys. I* (France) **4**, 863 (1994).
- [5] H. Takayasu, A.H. Sato, and M. Takayasu, *Phys. Rev. Lett.* **79**, 966 (1997).
- [6] E. Egener, T. Lux, and D. Stauffer, *Physica* (Amsterdam) **268A**, 250 (1999); D. Chowdhury and D. Stauffer, *Eur. Phys. J. B* **8**, 477 (1999).
- [7] R.N. Mantegna and H.E. Stanley, *Nature* (London) **376**, 46 (1995); Y. Liu *et al.*, *Physica* (Amsterdam) **245A**, 437 (1997); P. Cizeau *et al.*, *Physica* (Amsterdam) **245A**, 441 (1997); Y. Liu, P. Gopikrishnan, P. Cizeau, M. Meyer, C.-K. Peng, and H.E. Stanley, *Phys. Rev. E* (to be published); M.M. Dacorogna, U. A. Müller, R. J. Nagler, R. B. Olsen, and O. V. Pictet, *J. Int. Money Fin.* **12**, 413 (1993); L. Kullmann and J. Kertész, *Physica* (Amsterdam) **269A**, 98 (1999).
- [8] T. Lux, *Appl. Fin. Econ.* **6**, 463 (1996); P. Gopikrishnan, M. Meyer, L.A.N. Amaral, and H.E. Stanley, *Eur. Phys. J. B* **3**, 139 (1998); P. Gopikrishnan, V. Plerou, L.A.N. Amaral, M. Meyer, and H.E. Stanley, *cond-mat/9905305*; V. Plerou, P. Gopikrishnan, L.A.N. Amaral, M. Meyer, and H.E. Stanley, *cond-mat/9907161*; U. A. Müller, M.M. Dacorogna, R.B. Olsen, O. V. Pictet, M. Schwarz, and C. Morgengegg, *J. Bank. Fin.* **14**, 1189 (1990).
- [9] M.H.R. Stanley, L.A.N. Amaral, S.V. Buldyrev, S. Havlin, H. Leschhorn, P. Maass, M.A. Salinger, and H.E. Stanley, *Nature* (London) **379**, 804 (1996); H. Takayasu and K. Okuyama, *Fractals* **6**, 67 (1998); L.A.N. Amaral, S.V. Buldyrev, S. Havlin, M.A. Salinger, and H.E. Stanley, *Phys. Rev. Lett.* **80**, 1385 (1998); Y. Lee, L.A.N. Amaral, D. Canning, M. Meyer, and H.E. Stanley, *Phys. Rev. Lett.* **81**, 3275 (1998); V. Plerou, L.A.N. Amaral, M. Meyer, P. Gopikrishnan, and H.E. Stanley, *Nature* (London) **400**, 433 (1999).
- [10] A. Johansen and D. Sornette, *Eur. Phys. J. B* **9**, 167 (1999); *Physica* (Amsterdam) **261A**, 581 (1998); A. Arnoedo, J.-F. Muzy, and D. Sornette, *Eur. Phys. J. B* **2**, 277 (1998).
- [11] L. Laloux *et al.*, *Phys. Rev. Lett.* **83**, 1467 (1999); S. Galluccio *et al.*, *Physica* (Amsterdam) **259A**, 449 (1998).
- [12] R.N. Mantegna, *cond-mat/9802256*.
- [13] M. Levy, H. Levy, and S. Solomon, *Econ. Lett.* **45**, 103 (1994).
- [14] S. Ghashghaie *et al.*, *Nature* (London) **381**, 767 (1996); R.N. Mantegna and H.E. Stanley, *Nature* (London) **383**, 587 (1996); *Physica* (Amsterdam) **239A**, 255 (1997).
- [15] N. Vandewalle and M. Ausloos, *Int. J. Mod. Phys. C* **9**, 711 (1998); *Eur. Phys. J. B* **4**, 257 (1998).
- [16] E.P. Wigner, *Ann. Math.* **53**, 36 (1951).
- [17] T. Guhr, A. Müller-Groeling, and H.A. Weidenmüller, *Phys. Rep.* **299**, 190 (1998); M.L. Mehta, *Random Matrices* (Academic Press, Boston, 1991).
- [18] T. A. Brody *et al.*, *Rev. Mod. Phys.* **53**, 385 (1981).
- [19] The trades and quotes (TAQ) database. This database is published by the New York Stock Exchange and consists of 24 CD-ROMS for the period 1994–1995.
- [20] A. M. Sengupta and P. P. Mitra, *cond-mat/9709283*.
- [21] H. Bruus and J.-C. Anglès d’Auriac, *Europhys. Lett.* **35**, 321 (1996).
- [22] The number variance is defined as $\Sigma^2(L) \equiv \langle [N(\lambda + \frac{L}{2}) - N(\lambda - \frac{L}{2}) - L]^2 \rangle_\lambda$, where $N(\lambda) \equiv \sum_i \theta(\lambda - \lambda_i)$ is the integrated density of eigenvalues and $\langle \dots \rangle_\lambda$ denotes an average over λ [17,21].
- [23] The spectral rigidity is defined as $\Delta(L) \equiv \frac{1}{L} \langle \min_{A,B} \int_{\lambda-L/2}^{\lambda+L/2} [N(\lambda_1) - A\lambda_1 - B]^2 d\lambda_1 \rangle_\lambda$, where $\langle \dots \rangle_\lambda$ denotes an average over λ and $N(\lambda) \equiv \sum_i \theta(\lambda - \lambda_i)$ is the integrated density of eigenvalues [17,21].
- [24] Y.V. Fyodorov and A.D. Mirlin, *Phys. Rev. Lett.* **69**, 1093 (1992); **71**, 412 (1993); A.D. Mirlin and Y.V. Fyodorov, *J. Phys. A* **26**, L551 (1993).
- [25] P. A. Lee and T. V. Ramakrishnan, *Rev. Mod. Phys.* **57**, 287 (1985).
- [26] The large values of I_k for small eigenvalues indicate that the distribution of eigenvector components for the eigenvalues at the lower edge of the spectrum deviate from Gaussian prediction.
- [27] Metals or semiconductors with impurities can be described by Hamiltonians with random-hopping integrals [F. Wegner and R. Oppermann, *Z. Phys. B* **34**, 327 (1979)]. Electron-hopping between neighboring sites is more probable than hopping over large distances, leading to a Hamiltonian that is a random band matrix.
- [28] A random band matrix \mathbf{B} has elements B_{ij} independently drawn from different probability distributions. These distributions are often taken to be Gaussian and to be parametrized by their variance, which depends on i and j . Although such matrices are random, they still contain probabilistic information regarding the fact that a metric can be defined on their set of indices i .
- [29] A related idea for a hierarchical structure of the financial cross-correlation matrix was recently put forward in Ref. [12].